

Complex Energy Method for Scattering Processes

H. Kamada^{1) *)}, Y. Koike^{2,3) **)}, W. Glöckle^{4)***)}

¹⁾ *Department of Physics, Faculty of Engineering, Kyushu Institute of Technology,
Kitakyushu 804-8550, Japan*

²⁾ *Science Research Center, Hosei University, 2-17-1, Fujimi, Chiyoda-ku,
Tokyo 102, Japan*

³⁾ *Center for Nuclear Study, University of Tokyo, 2-1 Hirosawa, Wako,
Saitama 351-0198, Japan*

⁴⁾ *Institut für Theoretische Physik II, Ruhr-Universität Bochum,
D-44780 Bochum, Germany*

(Received February 5, 2008)

A method for solving few-body scattering equations is proposed and examined. The solution of the scattering equations at complex energies is analytically continued to get scattering T-matrix with real positive energy. Numerical examples document that the method works well for two-nucleon scattering and three-nucleon scattering, if the set of complex energies is properly chosen.

§1. Introduction

Scattering by more than two particles upon each other requires boundary conditions in configuration space of increasing complexity with growing particle numbers. Those boundary conditions are reflected in momentum space by $\epsilon \rightarrow 0$ limits of different resolvent operators $G(E + i\epsilon)$, where E is real and above some thresholds. While such a singularity can easily be handled on a two-body fragmentation cut, it causes already severe difficulties on a three-body fragmentation cut like in the three-body problem. There it leads to so called moving singularities^{1), 2)}. They can be treated directly for instance using spline interpolations or just straightforward subtraction methods³⁾. They can also be circumvented by contour deformations of the path of momentum integration^{4) - 7)}. In any case this requires great care. For more than three particles the four-body break up singularities pose even harder challenges.

In the past several additional methods have been proposed especially for two-body fragmentations. For instance in⁸⁾ the T-matrix is evaluated at negative energies, where bound state methods can be employed, and then analytically continued to the energy in the scattering region. In^{9) - 11)} the T-matrix is evaluated at several complex energies above the positive real energy axis and then the value on the real energy axis is again determined by analytical continuation. All the applications we are aware of are in the field of atomic physics.

The Lorentz transformation method used intensively by the Trento group¹²⁾

*) email: kamada@mns.kyutech.ac.jp

**) email: koike@i.hosei.ac.jp

***) email: walter.gloeckle@tp2.ruhr-uni-bochum.de

solves many body equations based on a complex energy, thereby avoiding the complicated boundary conditions in configuration space or when used in momentum space¹³⁾ avoiding singularities. But in this case no continuation to the real energy axis is performed and the imaginary part of the energy is kept fixed. Certain inversion procedures are subsequently employed.

Here we would like to reconsider the approach evaluating the two- and three-body T-matrix for complex energies and performing an analytical continuation to the real axis. In view of future applications to the field of nuclear physics we shall employ nucleon - nucleon forces. Section 2 displays our studies for two nucleons. We add a brief example for three-body scattering in section 3 end with a brief outlook in section 4.

§2. Two- Nucleon Scattering

We start with a simple model for the NN forces allowing for an analytical solution, the Yamaguchi separable force¹⁴⁾

$$v(p, p') = \lambda \frac{1}{p^2 + \beta^2} \frac{1}{p'^2 + \beta^2} \quad (2.1)$$

It leads to the two-body T- matrix

$$t(p, p'; E) = \tau(E) \frac{1}{p^2 + \beta^2} \frac{1}{p'^2 + \beta^2} \quad (2.2)$$

with

$$\tau(E) = (\lambda^{-1} + \frac{\pi(\beta^2 - k^2)m}{4\beta(\beta^2 + k^2)^2} + i\frac{\pi km}{2(\beta^2 + k^2)^2})^{-1} \quad (2.3)$$

Now we evaluate $\tau(E)$ replacing energy $E \equiv k^2/m$ by $E + i\epsilon$, where ϵ is now finite and we choose a set of discrete ϵ 's. That function $\tilde{\tau}(\epsilon) \equiv \tau(E + i\epsilon)$ can be analytically continued to $\epsilon = 0$. Padé approximant is well known as a method for the analytical continuation. It is useful when coefficients of the Taylor series are given. On the contrary, the functional values at several points are given in the present case. We employ the point method⁸⁾ and consider the continued fraction

$$\begin{aligned} \tilde{\tau}(\epsilon) &= \frac{\tau(E + \epsilon_1 i)}{1 + \frac{a_1(\epsilon - \epsilon_1)}{1 + \dots}} \\ &= \frac{\tau(E + \epsilon_1 i)}{1 +} \frac{a_1(\epsilon - \epsilon_1)}{1 +} \frac{a_2(\epsilon - \epsilon_2)}{1 +} \dots \end{aligned} \quad (2.4)$$

The coefficients a_i are easily deduced from the discrete set of $\tilde{\tau}(\epsilon)$ -values:

$$\begin{aligned} a_l &= \frac{1}{\epsilon_l - \epsilon_{l+1}} \left\{ 1 + \frac{a_{l-1}(\epsilon_{l+1} - \epsilon_{l-1})}{1 +} \frac{a_{l-2}(\epsilon_{l+1} - \epsilon_{l-2})}{1 +} \dots \right. \\ &\quad \left. + \frac{a_1(\epsilon_{l+1} - \epsilon_1)}{1 - [\tau(E + i\epsilon_1)/\tau(E + i\epsilon_{l+1})]} \right\} \end{aligned} \quad (2.5)$$

ϵ [MeV]	$\tau(E + i\epsilon)$ [fm ²]
0.1	-0.11701981630329802 -0.60857451562152798 i
0.2	-0.11924707235802381 -0.61103993137637069 i
0.3	-0.12151390252710244 -0.61348507952563247 i
0.4	-0.1238200220019596 -0.6159089815519192 i
0.5	-0.12616511343686601 -0.61831066360816722 i
0.6	-0.12854882700065837 -0.62068915788211088 i
0.7	-0.13097078049779975 -0.62304350396337027 i
0.8	-0.13343055955913999 -0.6253727502090104 i
0.9	-0.13592771790246552 -0.62767595510338259 i
1.0	-0.13846177766262588 -0.62995218860804847 i
0	-0.1148323866513037 -0.60608981409107898 i
exact	-0.11483238665130231 -0.60608981409107832 i

Table I. The propagator τ from Eq(2.3) for different ϵ - values. The value for $\epsilon=0$ is evaluated by the point method⁸⁾. The exact value is obtained from Eq.(2.3). The parameters are $\lambda=-0.5592[\text{fm}^2]$, $\beta=1.13[\text{fm}^{-1}]$ and $m^{-1}=41.47[\text{MeVfm}^2]$. The energy is 10.0 MeV.

ϵ [MeV]	T_{00} [fm ²]	T_{02} [fm ²]	T_{22} [fm ²]
0.4	0.074878-0.38063 i	$-9.7683 \times 10^{-4} + 8.8418 \times 10^{-3} i$	$4.5582 \times 10^{-3} - 2.9901 \times 10^{-4} i$
0.5	0.072896-0.38404 i	$-8.0207 \times 10^{-4} + 9.1311 \times 10^{-3} i$	$4.5523 \times 10^{-3} - 3.1989 \times 10^{-4} i$
0.6	0.070821-0.38742 i	$-6.2122 \times 10^{-4} + 9.4171 \times 10^{-3} i$	$4.5460 \times 10^{-3} - 3.4052 \times 10^{-4} i$
0.7	0.068651-0.39077 i	$-4.3437 \times 10^{-4} + 9.6996 \times 10^{-3} i$	$4.5392 \times 10^{-3} - 3.6090 \times 10^{-4} i$
0.8	0.066387-0.39409 i	$-2.4160 \times 10^{-4} + 9.9782 \times 10^{-3} i$	$4.5319 \times 10^{-3} - 3.8100 \times 10^{-4} i$
0	0.081860-0.36687 i	$-1.6137 \times 10^{-3} + 7.6562 \times 10^{-3} i$	$4.5768 \times 10^{-3} - 2.1337 \times 10^{-4} i$
exact	0.081860-0.36687 i	$-1.6138 \times 10^{-3} + 7.6564 \times 10^{-3} i$	$4.5767 \times 10^{-3} - 2.1338 \times 10^{-4} i$

Table II. The T-matrix for the 3S_1 - 3D_1 state and different ϵ -values. The value for $\epsilon=0$ is calculated by the point method and compared to the exact number resulting directly from Eq.(2.7).

and

$$a_1 = \frac{[\tau(E + i\epsilon_1)/\tau(E + i\epsilon_2)] - 1}{\epsilon_2 - \epsilon_1}. \quad (2.6)$$

An example is displayed in Table I, which shows the smooth behavior of $\tilde{\tau}(\epsilon)$ sampled for discrete ϵ - values according to Eq(2.3). The analytically continued value for $\epsilon=0$ as given by (2.4) agrees perfectly well with the exact one.

Now that approach also works perfectly for a modern high precision NN force like the CD-Bonn potential¹⁵⁾. As an example we take the coupled channel case $^3S_1 - ^3D_1$ and solve now the two- body Lippmann Schwinger equation

$$T_{l'l}(p', p) = V_{l'l}(p', p) + \sum_{l''} \int_0^\infty dp'' p''^2 V_{l'l''}(p', p'') \frac{1}{E + i\epsilon - \frac{p''^2}{m}} T_{l''l}(p'', p) \quad (2.7)$$

numerically for a set of ϵ - values. We use the same method. The continued fraction is now in the matrix

$$\tilde{T}(\epsilon) \equiv T(p, p; E + i\epsilon) = \frac{T(p, p; E + \epsilon_1 i)}{1 +} \frac{A_1(\epsilon - \epsilon_1)}{1 +} \frac{A_2(\epsilon - \epsilon_2)}{1 +} \dots \quad (2.8)$$

where A_i are components of 2×2 matrices. We arrive at the results displayed in Table II. There we show the on- shell T- matrix elements $T_{00}(p, p)$, $T_{02}(p, p)$, $T_{22}(p, p)$ for

$p = \sqrt{mE}$, $E = 10.0$ MeV and different ϵ -values. Again we see a smooth dependence on ϵ . The resulting matrix from the point method elements ($\epsilon=0$) are compared to the values evaluated by a standard subtraction method in Table II. The agreement is again perfect.

That expansion works equally well for off-shell T- matrices $\lim_{\epsilon \rightarrow 0} T(p', p; E + i\epsilon)$ where $p \neq p'$ and E not connected to p or p' .

§3. An example for three-nucleon scattering

We now proceed to three-nucleon scattering. Several methods have been developed to treat singularities for the Faddeev equations. The contour deformation method has been used as one of the methods for avoiding complicated singularities. Here we adopt a separable expansion method^{19), 20)} for calculating the two-body t-matrix. The resulting three-body equations become coupled integral equations in one variable and the contour deformation can be used as a tool for avoiding singularities. We compare the three-nucleon scattering amplitudes obtained by the present method with those from the traditional contour deformation method.

The three-body equations we solve are of the following form

$$X_{cc'}^{(J^\pi)}(q, q'; E) = Z_{cc'}^{(J^\pi)}(q, q'; E) + \sum_{c''} \int_0^\infty q''^2 dq'' Z_{cc''}^{(J^\pi)}(q, q''; E) \tau(E - q''^2/2\mu) X_{c''c'}^{(J^\pi)}(q'', q'; E) \quad (3.1)$$

Eqs. (3.1) are a generalization of the Amado-Mitra-Faddeev-Lovelace (AMFL) equations, which are three-body equations based on a simple rank-1 separable potential. The AMFL equations are described in standard textbooks for the quantum mechanical three-body problem and the generalization thereof in ref²¹⁾.

The three-body amplitude describing elastic scattering is the on-shell n-d amplitude $X_{c_i c_j}^{(J^\pi)}(q_0, q_0; E)$ with $E = q_0^2/2\mu$ and n-d channels c_i, c_j allowed for the total spin and parity J^π . The number of n-d channels are two for $J^\pi = (1/2)^+, (1/2)^-$ or three for others. The contour deformation utilizes the analytical continuation in the momenta q, q', q'' , while the present method is based on the analytical continuation in the energy E . In both cases, the various singularities in Eq.(3.1) are avoided.

We solve equ(3.1) with the present method at 10 MeV laboratory energy. At this energy, it is known that the breakup cross section is already about 10-20 % of the total cross section. Thus the moving singularity is quite important.

In Table III, we show a result for the three-nucleon scattering amplitude for $J^\pi = (1/2)^+$ based on the AV14 potential²²⁾. Two channels, denoted by d and q and reflecting spin states, are coupled for the n-d elastic scattering amplitude. The energy is chosen as $E = E_0 + i\epsilon$, where $E_0 = q_0^2/2\mu$ and q_0 is the initial momentum. In contrast to the simpler two-body singularity treated in section 2, where real epsilons were chosen, we use now also complex epsilons. This allows to have more complex energy points in the neighborhood of E_0 . Also we kept the imaginary parts of the energy always larger than 1 MeV (this is in relation to the chosen $E_0=10$ MeV) which guarantees a relatively smooth integrand and avoids problems in the

$i\epsilon$ [MeV]	$X_{dd}^{(J^\pi)}$ [fm ⁵]	$X_{dq}^{(J^\pi)}$ [fm ⁵]	$X_{qq}^{(J^\pi)}$ [fm ⁵]
0+1.0 i	0.5634 -1.7745 i	$-1.3755 \times 10^{-2} + 1.4227 \times 10^{-2} i$	$0.2618 + 4.5351 \times 10^{-2} i$
1.0+1.5 i	0.6669 -1.8203 i	$-1.9726 \times 10^{-2} + 8.4048 \times 10^{-3} i$	$0.2451 + 8.7036 \times 10^{-2} i$
0.0+2.0 i	0.7608 -1.8267 i	$-2.5101 \times 10^{-2} + 2.3236 \times 10^{-3} i$	$0.2198 + 0.1208 i$
0.5+1.5 i	0.6513 -1.9441 i	$-2.5979 \times 10^{-2} + 1.4182 \times 10^{-2} i$	$0.2878 + 0.1136 i$
-0.5+1.5 i	0.6937 -1.7363 i	$-1.4067 \times 10^{-2} + 2.8661 \times 10^{-3} i$	$0.2109 + 6.8861 \times 10^{-2} i$
0.5+1.0 i	0.4892 -1.9070 i	$-1.9834 \times 10^{-2} + 2.0626 \times 10^{-2} i$	$0.3135 + 5.9376 \times 10^{-2} i$
-0.5+1.0 i	0.6265 -1.6940 i	$-8.3854 \times 10^{-2} + 8.0987 \times 10^{-3} i$	$0.2225 + 3.6096 \times 10^{-2} i$
0.5+2.0 i	0.7819 -1.9252 i	$-3.1597 \times 10^{-2} + 7.5166 \times 10^{-3} i$	$0.2514 + 0.1543 i$
0	0.4248 -1.5440 i	$2.5154 \times 10^{-5} + 2.4663 \times 10^{-2} i$	$0.2586 - 4.5922 \times 10^{-2} i$
C. D.	0.4244 -1.5441 i	$4.9443 \times 10^{-5} + 2.4784 \times 10^{-2} i$	$0.2585 - 4.5875 \times 10^{-2} i$

Table III. T-matrix for three-body scattering for different complex ϵ -values and $J^\pi = (1/2)^+$. The value on the real axis ($i\epsilon=0$) is calculated by the point method and compared to the contour deformation number (C. D.) resulting directly from Eq.(3.1). The channels d and q correspond to spin doublet and quartet, respectively.

$i\epsilon$ [MeV]	$X_{dd}^{(J^\pi)}$ [fm ⁵]	$X_{dq}^{(J^\pi)}$ [fm ⁵]	$X_{qq}^{(J^\pi)}$ [fm ⁵]
0+1.0 i	0.2117 -0.3215 i	-0.2277 -0.2446 i	-0.6525 -0.6931 i
1.0+1.5 i	0.3032 -0.2590 i	-0.1941 -0.2410 i	-0.5818 -0.6991 i
0.0+2.0 i	0.3575 -0.1809 i	-0.1650 -0.2328 i	-0.5161 -0.6986 i
0.5+1.5 i	0.3999 -0.3379 i	-0.1851 -0.2748 i	-0.5806 -0.7702 i
-0.5+1.5 i	0.2454 -0.1941 i	-0.1984 -0.2116 i	-0.5775 -0.6340 i
0.5+1.0 i	0.2813 -0.4530 i	-0.2250 -0.2838 i	-0.6601 -0.7722 i
-0.5+1.0 i	0.1778 -0.2291 i	-0.2271 -0.2121 i	-0.6406 -0.6249 i
0.5+2.0 i	0.4549 -0.2156 i	-0.1526 -0.2606 i	-0.5111 -0.7635 i
0	$-4.5247 \times 10^{-2} - 0.3098 i$	-0.3013 -0.2316 i	-0.7976 -0.6440 i
C. D.	$-4.5393 \times 10^{-2} - 0.3098 i$	-0.3016 -0.2312 i	-0.7975 -0.6437 i

Table IV. Same as table III except for $J^\pi = (1/2)^-$.

convergence of the integration. As a consequence of that large imaginary parts of the discrete set of energies the amplitudes shown in Table 3 differ considerably from the amplitudes, C.D., on the real axis as obtained with the contour deformation method. Nevertheless the point method applied to those amplitudes leads to good converged results which are very close to the amplitudes C.D.

Table IV shows the amplitudes for $J^\pi = (1/2)^-$. We see again the same good convergence to the results achieved by the contour deformation method.

§4. Outlook

We demonstrated the easiness by which the singularities in the Lippmann Schwinger equation for NN scattering can be avoided. We used a set of complex energies above the positive real axis and based on that performed an analytical continuation to the real axis. This analytical continuation was performed by the point method (a continued fraction expansion).

In case of the three-body Faddeev equations one encounters in the integral kernel two types of singularities related to the nucleon-deuteron and three-nucleon cuts. The first one is generated by the off- shell NN T- matrix and the second one by the

free three-nucleon propagator. As shown in section 3 the complex energy method can again be applied. The resulting accuracy is quite satisfactory and the resulting amplitudes agree very well with the ones from the contour deformation method.

This method also allows to search for resonances below the real axis. In a forthcoming paper we will display that this method is applicable to even more than three particles like the complex four- body break- up process , where no calculations have been performed so far.

Acknowledgements

One of authors (H.K.) is grateful to I. Fachruddin, A. Hemmdan, E. Epelbaum, R. Skibiński and H. H. Oo not only for numerous discussions but also for the kind atmosphere when he recently visited the Ruhr-Universität Bochum.

-
- 1) E. W. Schmid and H. Ziegelmann, "The Quantum Mechanical Three-Body Problem"(Pergamon Press, Oxford 1974)
 - 2) D. Hüber, H. Kamada, H. Witała, W. Glöckle, Few-Body Systems **16**, 165 (1994).
 - 3) W. Glöckle, H. Witała, D. Hüber, H. Kamada, and J. Golak, Phys. Rep. **274**, 107 (1996).
 - 4) C. Lovelace, Phys. Rev C **135** , 1125 (1964).
 - 5) R. T. Cahill, I. H. Sloan, Nucl. Phys. **A165**, 161 (1971).
 - 6) W. Ebenhöh, Nucl. Phys. **A191**, 97 (1972).
 - 7) Y. Koike, Nucl. Phys. **A301**, 411 (1978).
 - 8) L. Schlessinger, Phys. Rev. **167**, 1411 (1968).
 - 9) F. A. McDonald, J.Nuttall, Phys. Rev. A **4**, 1821 (1971).
 - 10) W. P. Reinhardt, D. W. Oxtoby, T. N. Resicigno, Phys. Rev. Lett. **28**, 401 (1972).
 - 11) G. Doolen, G. McCartor, F. A. McDonald, J. Nuttall Phys. Rev A **4**, 108 (1971).
 - 12) V. D. Efros, W. Leidemann, G. Orlandini, Phys. Lett. B **338**, 130 (1994); V. D. Efros, Yad. Fiz. **41**, 1498 (1985) [Sov. J. Nucl. Phys. **41**, 949 (1985)].
 - 13) S. Martinelli, H. Kamada, G. Orlandini, and W. Glöckle, Phys. Rev. C **52**, 2906 (1995).
 - 14) Y. Yamaguchi, Phys. Rev **95** , 1628 (1954).
 - 15) R. Machleidt, F. Sammarruca, and Y. Song, Phy. Rev. C**53**, R1483 (1996).
 - 16) L. D. Faddeev, Sov. Phys. JETP **12** , 1014 (1961).
 - 17) E. O. Alt, P. Grassberger, W. Sandhas, Nucl. Phys. **B2** , 167 (1967).
 - 18) W. Glöckle, *The Quantum-Mechanical Few-Body Problem* (Springer Verlag, Berlin, Heiderberg, 1983).
 - 19) Y. Koike, Phys. Rev. C **42**, R2286 (1990).
 - 20) Y. Koike, W. C. Parke, L. C. Maxmon, and D.R. Lehman, Few-Body Systems **23**, 53 (1997).
 - 21) W. C. Parke, Y. Koike, D. R. Lehman, and L. C. Maxmon, Few-Body Systems **11**, 89 (1991).
 - 22) R. B. Wiringa, R. A. Smith, and T. L. Ainsworth, Phys. Rev. C **29**, 1207 (1984).